

Date of report 03 Feb 2020

Reported case interaction between Cobicistat and Ranolazine

Drugs suspected to be involved in the DDI

Perpetrator	Daily Dose
Cobicistat	150 (mg)
Dose adjustment performed No	Administration Route Oral
Start date	End date
Jan. 1, 2014	Ongoing
Victim	Daily Dose
Ranolazine	1000 (mg)
Dose adjustment performed No	Administration Route Oral
Start date	End date
Nov. 1, 2018	Jan. 31, 2019

Complete list of drugs taken by the patient

Antiretroviral treatment Darunavir (with Ritonavir or Cobicistat) Emtricitabine/Tenofovir-AF

Complete list of all comedications taken by the patient, included that involved in the DDI

Aspirin 81 mg once daily (5 years), cilostazol 50 once daily (5 years), clopidogrel 75 mg once daily (5 years), atorvastatin 40 mg once daily (5 years), tiotropium 18 ug once daily (2 years), ondansetron 8 mg twice daily (<2months), pantoprazole 40 mg once daily (<2months), tramadol 50 mg once daily (1 year), ranolazine 500 mg twice daily (2 months)

Clinical case description	
Gender	Age
Male	64
eGFR (mL/min)	Liver function impairment
>60	No

Description

In early January 2019, a 64 year old man was admitted to the hospital with left-sided chest pain, dizziness and near syncope. The patient also reported persistent and severe episodes of nausea, vomiting, dyspepsia and anorexia for the past 2 months. Electrocadiography revealed first-degree atrioventricular block (AV). During the hospitalization, the pharmacist identified the drug-drug interaction between ranolazine and darunavir/cobicistat which led to the interruption of ranolazine. The patient's chest pain, nausea and dizziness resolved in a couple of days after stopping ranolazine. Gastrointestinal and AV block events are known side effects of ranolazine. In the present case, these side effects occurred likely due to cobicistat inhibition of ranolazine metabolism leading to an increased exposure. This case has been published by Dougherty JA et al. Ann Pharmacother 2019; 53:966-7.

Clinical Outcome

Toxicity

Drug Interaction Probability Scale (DIPS)

Score

8 - Probable

Editorial Comment

Ranolazine levels have been shown to be significantly increased by ketoconazole another strong CYP4A4 inhibitor (Jerling M et al. J Clin Pharmacol 2005). Ranolazine product label contra-indicated the coadministration with strong CYP3A4 inhibitors.

University of Liverpool Recommendation

These drugs should not be coadministered

For more information click here