



Date of report 22 Jun 2020

Reported case interaction between **Cobicistat** and **Clorazepate**

Drugs suspected to be involved in the DDI

Perpetrator

Cobicistat

Daily Dose

150 (mg)

Dose adjustment performed

No

Administration Route

Oral

Start date

Dec. 3, 2019

End date

Ongoing

Victim

Clorazepate

Daily Dose

50 (mg)

Dose adjustment performed

No

Administration Route

Oral

Start date

Unknown

End date

Ongoing

Complete list of drugs taken by the patient

Antiretroviral treatment

Darunavir/Cobicistat/Emtricitabine/Tenofovir-AF

Complete list of all comedications taken by the patient, included that involved in the DDI

Clorazepate

Clinical case description

Gender

Female

Age

50

eGFR (mL/min)

>60

Liver function impairment

No

Description

50 years old woman diagnosed with HIV in 1997. She had been receiving ART with abacavir, lamivudine and nevirapine maintaining undetectable plasma viral load until September 2019 when she stopped ART during 2 months. In November 2019 ART was reinitiated with FTC/TAF + RAL but it was switched to DRV/c/FTC/TAF because of poor adherence. She was also receiving clorazepate because of a chronic anxiety disorder. Clorazepate is converted to nordiazepam which is then metabolized to oxazepam by CYP3A4. Despite nordiazepam plasma concentrations are expected to be increased in combination with a strong inhibitor of CYP3A4 like cobicistat, no adverse events were observed.

Clinical Outcome

No unwanted outcome

Editorial Comment

Darunavir/c is expected to increase nordiazepam exposure due to CYP3A4 Inhibition. Development of adverse effects as a consequence of drug-drug interaction is likely to vary from one person to the other.

University of Liverpool Recommendation

- Potential interaction - may require close monitoring, alteration of drug dosage or timing of administration

For more information [click here](#)