



Date of report 28 Jul 2020

Reported case interaction between **Etravirine** and **Fluconazole**

Drugs suspected to be involved in the DDI

Victim

Etravirine

Daily Dose

400 (mg)

Dose adjustment performed

No

Administration Route

Oral

Start date

Unknown

End date

Ongoing

Perpetrator

Fluconazole

Daily Dose

400 (mg)

Dose adjustment performed

No

Administration Route

Intravenous

Start date

April 12, 2020

End date

May 29, 2020

Complete list of drugs taken by the patient

Antiretroviral treatment

Darunavir (with Ritonavir or Cobicistat)
Etravirine

Complete list of all comedications taken by the patient, included that involved in the DDI

Fluconazole, atorvastatin, cloxacillin, enalapril, enoxaparin, metamizol, insulin, paracetamol, omeprazol

Clinical case description

Gender
Male

Age
54

eGFR (mL/min)
60-30

Liver function impairment
No

Description

This is a HIV patients with an acute infection of a hip prosthesis caused by a *Candida rugosa*, that was isolated in surgical cultures and treated with fluconazol. Fluconazol inhibits etravirine metabolism and coadministration may increase etravirine exposures (AUC by 86%, C_{max} by 75%, C_{min} by 2.09-fold) while it has no significant effect on fluconazole exposure. No dose adjustments were done. The patient presented a multifactorial severe anemia that may be worsened by an increased exposure to etravirine and that improved when fluconazole treatment was stopped.

Clinical Outcome

Toxicity

Drug Interaction Probability Scale (DIPS)

Score

3 - Possible

Editorial Comment

Coadministration significantly increased etravirine exposures (AUC by 86%, Cmax by 75%, Cmin by 2.09-fold) and had no significant effect on fluconazole exposure. Although no dose adjustments are needed, etravirine and fluconazole should be coadministered with caution as there are limited safety data on increased etravirine exposure. Anemia is among common laboratory abnormalities (4%) in etravirine prescribing information, although no clear relationship with etravirine concentrations has been described.

University of Liverpool Recommendation

■ Potential interaction - may require close monitoring, alteration of drug dosage or timing of administration

For more information [click here](#)