



Date of report 02 Sep 2021

## Reported case interaction between **Nevirapine** and **Warfarin**

### Drugs suspected to be involved in the DDI

Perpetrator

**Nevirapine**

Daily Dose

400 (mg)

Dose adjustment performed

No

Administration Route

Oral

Start date

Unknown

End date

Unknown

Victim

**Warfarin**

Daily Dose

8.75 (mg)

Dose adjustment performed

No

Administration Route

Oral

Start date

Unknown

End date

Unknown

## Complete list of drugs taken by the patient

Antiretroviral treatment

Nevirapine

Abacavir/Lamivudine

Complete list of all comedications taken by the patient, included that involved in the DDI

warfarin maintenance therapy on nevirapine treatment: 8.75 mg/d from Monday to Saturday and 7.5 mg on Sunday

## Clinical case description

Gender

Female

Age

50

eGFR (mL/min)

>60

Liver function impairment

No

Description

A 50-year-old woman HIV-infected was referred to evaluate the cause of uncontrolled INR. The patient had been on maintenance warfarin therapy for the treatment of deep venous thrombosis for more than 10 years. Under treatment with a nevirapine based regimen, warfarin maintenance dosage was 8.75 mg/day from Monday to Saturday and 7.5 mg on Sunday which allowed to achieve a stable therapeutic INR (between 1.72 and 2.37). On November 2020, the antiretroviral treatment was changed to dolutegravir/lamivudine for treatment simplification. A control of INR 20 days later showed a value of 2.29. In December, the patient noted gum bleeding but did not go to the hospital to measure the INR due to the second COVID-19 pandemic wave. The patient reduced the warfarin dose on her own to 7.5 mg/day

however the gingival bleeding persisted. An INR was finally measured on December 29, 2020 and was 4.19 confirming the over-anticoagulation. The antiretroviral treatment was changed back to the nevirapine containing regimen and INR was again in the therapeutic range. This case was published by Cattaneo D et al. in AIDS 2021.

## Clinical Outcome

### Toxicity

## Drug Interaction Probability Scale (DIPS)

Score

**8 - Probable**

## Editorial Comment

The observed interaction is explained by the fact that warfarin dosage had to be increased under nevirapine treatment to compensate for nevirapine inducing effect. Now, switching from nevirapine to dolutegravir resulted in an increased anti-coagulant effect given that warfarin dosage had not been adapted to account for the fact that dolutegravir has no inducing effect. This case illustrates the importance of re-adjusting the comedication dosage when switching from an antiretroviral drug with inhibitory/inducing properties to an antiretroviral drug devoid of effects on drug metabolizing enzymes. In case studies of three Italian patients all taking antiretroviral therapy including nevirapine

plus warfarin, all experienced increased Quick times. In the first case 5 mg of warfarin per day was able to stabilise anticoagulant activity. In the second case 12 mg of warfarin was required and in the third case only after stopping nevirapine was 7.5 mg of warfarin enough to achieve the therapeutic range needed. Need for increased dose of warfarin in HIV patients taking nevirapine. Dionisio D, Mininni S, Bartolozzi D et al. AIDS, 2001,15:277-78.

## University of Liverpool Recommendation

- Potential interaction - may require close monitoring, alteration of drug dosage or timing of administration

For more information [click here](#)